



***DECODING EARLY TRANSPLANT GRAFT BIOPSIES
- A TRANSCRIPTOMIC ANALYSIS OF PANCREAS EARLY
ACUTE REJECTION***

Charlotte Franken, Tim Swaab, Jordi Rovira, Ivan Archilla, M^a Angeles GarciaCriado, Joana Ferrer-Fàbrega, Fritz Diekmann, Maria Ramirez-Bajo, Pedro Ventura-Aguar

LENIT – Laboratori Experimental de Nefrologia i Trasplantament



Pancreas transplant acute rejection



Incidence of acute rejection (AR)

10 – 35 % in first 12 months post-Tx

Risk factors for rejection

- Transplant type: SPK < PAK < PTA
- Donor age
- DSA presence
- Previous AR episode


Pancreas transplant acute rejection

American Journal of Transplantation 2013; 13: 1019–1025
Wiley Periodicals Inc.

© Copyright 2013 The American Society of Transplantation
and the American Society of Transplant Surgeons

doi: 10.1111/ajt.12167

Acute Pancreas Allograft Rejection Is Associated With Increased Risk of Graft Failure in Pancreas Transplantation



Incidence of acute rejection (AR)


10 – 35 % in first 12 months post-Tx

Risk factors for rejection

- Transplant type: SPK < PAK < PTA
- Donor age
- DSA presence
- Previous AR episode


➔ AR is associated with increased risk of graft failure

Pancreas graft rejection analysis




➔ **Need:** Development of **Pancreas-specific** genesets

Study design




Methods

- Samples: FFPE pancreas biopsies taken up to 60 days post-Tx (n = 20)
- B-HOT panel: 770 genes associated with graft injury




Workflow




Data analysis




Pancreas biopsy



FFPE samples



nanostRING



Study population characterisation

	Overall (n=20)	No rejection (n= 7)	Rejection (n= 13)	P value
				
Recipient age (years; mean±SD)	39±8,3	43,1±9	36,8±7,4	0,11
Donor age (years; median (IQR))	33,5 (21,3 – 45,0)	36 (19,0 – 46,0)	31 (22,5 – 45,0)	0,84
Time between PTx and Bx (days, mean±SD)	28,6±14,6	28,7±10,8	28,5±16,6	0,97
Transplant type (n, %): SPK PAK	16 (80%) 4 (20%)	6 (85,7%) 1 (14,3%)	10 (76,9%) 3 (23,1%)	1,00
cPRA (%; median (IQR))	0 (0,0 – 50,3)	0 (0,0 - 51)	0 (0 – 49,5)	0,86
Total HLA mismatches (median (IQR))	5 (4 – 6)	6 (5 – 6)	5 (4 – 5)	0,15
CIT (h, median (IQR))	8,5 (7,1 – 10,5)	9 (7,9 – 12,0)	8 (6,9 – 10,4)	0,21

*All recipients received standard induction therapy, consisting of tacrolimus, prednisone, and mycophenolate mofetil (MMF)

Histological characterisation



	Overall (n=20)	No rejection (n= 7)	Rejection (n= 13)	P value
Septal inflammation (n, %): 1+	14 (70%)	3 (42,9%)	11 (84,6%)	0,03
2+	1 (5%)	0 (0%)	1 (7,7%)	
Ductal inflammation (n, %): 1+	11 (55%)	0 (0%)	11 (84,6%)	< 0,001
2+	1 (5%)	0 (0%)	1 (7,7%)	
Acinar inflammation (n, %): 1+	6 (30%)	1 (14,3%)	5 (38,5%)	0,43
2+	1 (5%)	0 (0%)	1 (7,7%)	
Arteritis (n, %): +	1 (7,1%)	0 (0%)	1 (10%)	1,00
Venulitis (n, %): +	7 (35%)	0 (0%)	7 (53,8%)	0,04
Isletitis (n, %): 1+	2 (10%)	0 (0%)	2 (15,4%)	0,68
2+	1 (5%)	0 (0%)	1 (7,7%)	
C4d: +	1 (5%)	0 (0%)	1 (7,7%)	1,00
Fibrosis: +	2 (10,5%)	1 (14,3%)	1 (8,3%)	1,00

Histological characterisation – IHQ


No rejection

CD3


CD68




Rejection



T cell marker




Macrophage marker




Acute Rejection – Differently expressed genes

117 significantly differently expressed genes → 31 underexpressed & 86 overexpressed


Rejection vs No Rejection (baseline)




Acute Rejection – Cell type scores¹




Cytotoxic cell score




Exhausted CD8 cell score




Macrophage score




Dendritic cell score




CD45 score




T-cell score



CD8 T-cell score




IgM PC score





¹ cell type scores calculated based on the median expression of cell-specific genes

Acute Rejection – Pathway signature scores¹




¹ pathway signature scores calculated based on the median expression of pathway-specific genes

Correlation histology & cell type scores



Graft survival


	Overall (n=20)	No rejection (n= 7)	Rejection (n= 13)	P value
Pancreas graft survival (n, %): Failed	5 (25%)	0 (0,0%)	5 (38,5%)	0,117
Functioning	15 (75%)	7 (100%)	8 (61,5%)	



Graft survival – Differently expressed genes

88 significantly differently expressed genes → 45 underexpressed & 43 overexpressed

Graft failure vs Graft survival (baseline)



45 genes ↓*

- MARCH8: NF-kappaB signaling
- AKR1C3: KT1, Oxidative stress
- DDX50: NF-kappa-B & IRF3 Signaling
- XBP1: Eosinophils, Oxidative stress
- HDAC3: Chromatin modifying enzymes, NOTCH signaling

43 genes ↑*

- EZH2: Chromatin modifying enzymes, Oxidative stress
 - PTX3: M2 transition
 - S100A9: M2 transition
 - TFRC: M2 transition
 - S100A8: M2 transition
- } M2

Conclusions

Transcriptomic analysis revealed distinct differences between the Rejection and No Rejection group

1. **Cell-type score comparisons** demonstrated significant differences in **dendritic cell** and **T-cell-related scores** between the rejection and no-rejection groups
2. **Pathway signature analysis** indicated significantly higher scores for **M2 macrophages**, **IL-7** and **IL-2** signaling, **Th17** and **Th2** differentiation, and **TCMR-related pathways** in the rejection group
3. Recipients with **graft failure** exhibited a significantly **higher expression** of **M2** macrophage-associated genes
4. A deeper understanding of the **molecular drivers** of acute pancreas transplant rejection could enhance **diagnostic accuracy** and support the development of a **pancreas-specific gene panel**, with potential therapeutic implications

Acknowledgements

LENIT – Laboratori Experimental de Nefrologia i Trasplantament

Fritz Diekmann
Pedro Ventura-Aguar
Maria Ramirez-Bajo
Elisenda Banon-Maneus
Jordi Rovira
Tim Swaab
Natalia Hierro-Garcia

